



Post-harvest fungal rot of green pepper fruits (*Capsicum annuum*): methanolic extracts of tropical spices as bio rescue agents

J. Y Ijato^{1*}, O. M. Obembe¹, O. K Salami¹, O. O Olajide², B. O Ojo³, H. O Yakubu¹ and B.A Adanikin⁴

¹Department of Plant Science and Biotechnology, Faculty of Science, Ekiti State University, P.M.B 5363, Ado-Ekiti, Ekiti State, Nigeria.

²Department of Crop, Horticulture and Landscape Design, Faculty of Agriculture, Ekiti State University, P.M.B 5363, Ado-Ekiti, Ekiti State, Nigeria.

³Department of Biology, The Polytechnic, Ibadan, Nigeria.

⁴Nigeria Agricultural Quarantine Service, Moore Plantation, Apata, Ibadan, Nigeria.

*Corresponding author: considerureternity@gmail.com

ABSTRACT: This study aimed at evaluating the control potential of some essential oils on green pepper fruit (*Capsicum annuum*) rot. Samples of green pepper fruits (infected and healthy) and spices were obtained from Ado-Ekiti market, fungal rot pathogens were isolated from infected green pepper fruit and pathogenicity test was carried out to authenticate the pathogenic status of the fungal isolates. The fungi isolated from the green pepper fruits were: *Aspergillus spp*, *Cladosporium spp*, *Fusarium solani*, *Rhizopus stolonifer*, *Mucor spp*, *Alternaria spp*. The fungal isolates were identified using cultural and morphological features such as colony growth pattern, conidial morphology and pigmentation. Extracts of clove, ginger, and garlic were obtained using established standards. The fungal inoculum was prepared from 5-days old culture grown on potato dextrose agar and the effects of the essential oils were determined on the pepper fruit rot. The sensitivity of the fungal isolates to the extracts was found by assessing the diameter of the zone of inhibition in which significant susceptibility was taken as 25 mm in diameter. It was observed that the mycelia growth inhibition value by *Capsicum annuum* essential oils at 10⁵ concentration were *Aspergillus spp* (32.85mm), *Cladosporium spp* (33.57) mm, *Fusarium solani* (33.12) mm, *Rhizopus stolonifer* (33.22) mm, *Mucor spp* (32.935mm) was on *Fusarium solani*.

[J. Y Ijato, O. M. Obembe, O. K Salami, O. O Olajide, B. O Ojo, H. O Yakubu and B.A Adanikin. **Post-harvest fungal rot of green pepper fruits (*Capsicum annuum*): methanolic extracts of tropical spices as bio rescue agents.** J Am Sci 2022;18(9):1-8]. ISSN 1545-1003 (print); ISSN 2375-7264 (online). <http://www.jofamericanscience.org>. 01. doi:10.7537/marsjas180922.01.

Keywords: Bio agents, green pepper, post-harvest, tropical spices.

INTRODUCTION

Peppers are native to Mexico, Central America, and northern South America. Pepper seeds were imported to Spain in 1493 and then spread through Europe and Asia. The mild bell pepper cultivar was developed in the 1920s, in Szeged, Hungary (Sasvari, Joanne, 2005). Pepper (*Capsicum annuum*) is a genus of flowering plants in the nightshade family, Solanaceae. *Pepper Capsicum* consists of approximately 20-27 species out of which *Capsicum annuum*, *Capsicum baccatum*, *Capsicum chinense*, *Capsicum frutescens* and *Capsicum pubescens* were domesticated (Mason, 2010). In low income Sub-Saharan African countries, the incidences of post-harvest losses (from farm to fork) are the main causes of food insecurity (Zorya *et al.*, 2011). Rot fungi are ubiquitous biological agents that are able to infect fruits because of their ability to produce a wide range of hydrolytic enzymes. Mould infection depends on many factors such as pH, water activity, temperature,

atmosphere, time, etc (Magan and Aldred, 2007). Fungal infection can generate quality loss and health hazards through the production of toxins (Parveen *et al.*, 2016).

Moreover, pathogenic infection in the field or in post-harvest storage can affect the health of humans and livestock, especially if the pathogen produces toxins in or on consumable products (Menzler-Hokkanen, 2006). These reduce their yield and market values before and after harvest as fungi make way into plant host tissue through natural openings such as lenticels, stomata and through the unbroken epidermis by means of appressorium or germ tube (Ademoh *et al.* 2017). Various methods, strategies, and approaches are used in the management of plant diseases. These approaches have contributed significantly to the remarkable improvements in crop productivity and quality of crops produced (Chandrashekhara *et al.*, 2012). Therefore, there is a need to isolate and identify

fungi associated with post-harvest rot in pepper (*Capsicum annum*) in order to control the pathogens.

MATERIALS AND METHODS

Collection of plant materials

Three plants (*Syzygium aromaticum*, *Zingiber officinale* and *Allium sativum*) used in this study were collected from a garden at bank road, Ado Ekiti. Identity of the plant samples was authenticated using the herbarium specimens of the Department of Plant Science and Biotechnology, Ekiti State University, Ado Ekiti, Nigeria.

Preparation of plant extracts

The plant samples were washed in tap water, cut into small pieces of about 2 mm x 2 mm in size. The sample pieces were air dried for 15 days under room temperature (28°C). These were separately pulverized using an electric blender (Model M 20 IKA Universal Mill, IKA Group Japan). About 1000g each of the finely ground samples were carefully weighed and separately soaked in 1000 ml of 95 % methanol. The plant materials were soaked for 7 days. The supernatant was decanted into clean labeled flask and adequately corked. About 850 ml of methanol was further added to each of the plant samples left to stand for 3 days.

The resulting supernatant solution was separately decanted and added to the first, the crude extracts were filtered with sterile filter paper into labeled conical flask. The filtrates were concentrated in a rotary evaporator (Optional Oil Bath Model RE-3020 America (ambient to 180°C). The plant extract obtained was transferred into a labeled beaker and allowed to stand at room temperature for 24 hours to permit evaporation of the residual solvent. The extracts were kept in a refrigerator at 4°C until used (Fagbohun and Bamikole, 2019).

Preparation of media

The media used were prepared, sterilized and used according to manufacturer's instruction.

Isolation and identification of the fungal organisms

Diseased portion of the pepper fruits were cut under aseptic conditions into small bits of 5mm into a sterile dish with the aid of scissors which was flamed over a Bunsen's burner flame and dipped inside methylated spirit (Fawole and Oso, 1995, Ijato *et al*, 2022 a & b). The cut diseased bits sterilized with 70% ethanol were then placed centrally on Petri dishes containing solidified potato dextrose agar (PDA). The solidified plates were incubated at room temperature (28 + 2°C) in the dark for 72h. The fungal colonies growing from the incubated plates were sub-cultured into fresh medium until pure culture was obtained.

Microscopic examination was used after examining the colony characteristics to establish identity of fungi.

A sterile needle was used in taking a little portion of the hyphae containing spores and placed on the sterile glass slide and then stained with lactophenol cotton blue and examined under the microscope for fungal structures. The morphology and culture characteristics observed were compared with structures in (Snowdon, 1990). Forty grams (40g) PDA powder was placed in five liters conical flask. One hundred millilitre (100 mL) distilled water was added and boiled to completely dissolve the powder. To prevent bacterial growth, 0.2g of streptomycin was added to the potato dextrose broth. The supernatant was carefully transferred into sterile conical flasks and autoclaved at 120°C for 15 minutes at 101bf pressure and poured into Petri dishes for solidification.

Identification of isolates

Morphology of the fungal isolates was macroscopically studied by observing the colony features (color, shape and size) (Cheesbrough, 2000), and microscopically studied was by staining slide mounted with a small portion of the mycelium with a lactophenol cotton blue and examined under the compound microscope using x40 objective (Sohail and Bayan, 2018).

Pathogenicity test

The approach of Balogun *et al.*, (2005) was employed to determine the pathogenicity of the various fungal isolates. Apparently healthy and matured pepper fruits were surfaced-sterilized with 0.5% sodium hypochlorite for 30 seconds and then rinsed in three changes of sterile distilled water. A 5 mm diameter flame-sterilized cork borer cylindrical cores were removed from each fruit and were then inoculated aseptically with 5 mm diameter disc from the advancing edge of 7-day-old fungal culture of any one isolate.

Petroleum jelly was smeared to completely seal the surface of each of the inoculated pepper fruit to prevent external infection before incubating for 10 days in three replicates. The controls were inoculated with a disc of solidified potato dextrose agar medium. Fruits were inoculated in three replicates. Rot symptoms developed with different fungal isolates were compared to the natural original rot. The pathogens were re-isolated and identified using the same procedures described earlier in this finding.

Determination of antifungal activities of test plants

About 1 mL each of varying concentrations viz: 0.25, 0.50, 0.75 and 1.0 % of crude extracts were dispensed separately onto the 15 mL molten medium in Petri dishes with the help of a syringe. These were

thoroughly mixed and allowed to solidify. A portion of fungal growth from the periphery was cut with the help of a sterilized cork borer of 6 mm in diameter and placed onto the middle on each Petri dish and incubated at room temperature. A week after incubation, measurement of the colony was taken directly with the help of a scale. Control experiment was set up without adding any plant extract.

Difference in the measurement between the mycelia growth of the fungi and that of the cork borer gives the growth of colony and the effect of different concentration of extraction the growth of the species. Fungi toxicity was recorded in terms of percentage mycelia growth inhibition which was calculated using the formula as described by Nene and Thapilyal (2002), Ijato *et al.*, 2022 a & b

$$I = \frac{DC - DT}{DC} \times 100$$

Where: I = Percentage inhibition, DC = Average diameter of control (8cm), DT = Average diameter of growth with treatment.

Proximate analysis

The proximate composition was determined according to AOAC (2000).

Moisture content determination

Two gram (2.0g) of each sample was placed in an oven maintained at 100 -103°C for 16 hours with the weight of the wet sample and the weight after drying noted. The drying was repeated until a constant weight was obtained. The moisture content was expressed in terms of loss in weight of the wet sample.

$$\% \text{ moisture content} = \frac{\text{weight of moisture}}{\text{weight of sample}} \times 100$$

Ash content determination

Two gram (2.0g) of each of the oven-dried samples in powder form were accurately weighed and placed in crucible of known weight. These were ignited in a muffle furnace and ashed for 8 hours at 550°C. The crucible containing the ash was then removed, cooled in a desiccator and weighed and the ash content expressed in term of the oven-dried weight of the sample.

$$\% \text{ Ash content} = \frac{\text{weight of ash}}{\text{weight of sample}} \times 100$$

Protein content determination

The protein nitrogen in 1g of the dried samples was converted to ammonium sulphate by digesting with concentrated H₂SO₄ in the presence of CuSO₄ and Na₂SO₄. These were heated and the

ammonia evolved was steam distilled into boric acid solution. The nitrogen from ammonia was deduced from the titration of the trapped ammonia with 0.1M HCl with Tashirus indicator (double indicator) until a purplish pink color was obtained. Crude protein was calculated by multiplying the value of the deduced nitrogen by the factor 6.25mg.

Crude fiber content determination

A 2.0g of each sample was weighed into separate beakers, the samples were then extracted with petroleum ether by stirring, settling and decanting 3 times. The samples were then air dried and transferred into a dried 100ml conical flask. 200cm³ of 0.127M sulphuric acid solution was added at room temperature to the samples. The first 40cm³ of the acid was used to disperse the sample. This was heated gently to boiling point and boiled for 30 minutes. The contents were filtered to remove insoluble materials, which was then washed with distilled water, then with 1% HCl, next with twice ethanol and finally with diethyl ether. Finally, the oven-dried residue was ignited in a furnace at 550°C. The fiber contents were measured by the weight left after ignition and were expressed in term of the weight of the sample before ignition.

Fat content determination

The lipid content was determined by extracting the fat from 10g of the samples using petroleum ether in a Soxhlet apparatus. The weight of the lipid obtained after evaporating off the petroleum ether from the extract gave the weight of the crude fat in the sample.

Carbohydrate content determination

The carbohydrate content of the samples were determined as the difference obtained after subtracting the values of protein, lipid, ash and fiber from the total dry matter (AOAC 2009).

Data analysis

Data obtained were subjected to one way analysis of variance while the means were compared by Duncan's New Multiple Range Test at 95 % confidence interval using Statistical Package for Social Sciences version. Differences were considered significant at p≤0.05.

RESULTS

Table 1 shows the fungi isolated from the stored green pepper fruits: *Aspergillus spp*, *Cladosprium spp.*, *Fusarium solani*, *Rhizopus stolonifer*, *Mucor spp* and *Alternaria* and the frequency of occurrence of fungal isolate from green pepper fruits, it was observed that *Aspergillus spp* had the highest occurrence of (36%), followed by

Alternaria spp., (28%), *Cladosporium spp.*, (16%), *stolonifer* (4%),
Fusarium solani (8%), *Mucor spp* (8%) and *Rhizopus*

Table 1. Percentage frequency of fungi isolates from stored green pepper fruits.

| Fungi species | Number of isolates | Frequency (%) |
|----------------------------|--------------------|---------------|
| <i>Aspergillus spp</i> | 21 | 29.6 |
| <i>Cladosporium spp</i> | 14 | 19.7 |
| <i>Fusarium solani</i> | 8 | 11.3 |
| <i>Rhizopus stolonifer</i> | 17 | 23.9 |
| <i>Mucor spp</i> | 2 | 2.8 |
| <i>Alternaria spp</i> | 9 | 12.7 |
| Total | 71 | 100 |

The antifungal effects of methanol extract of *Syzygium aromaticum*, *zingiber officinale* and *Allium sativum* on fungal isolate from rotten green pepper fruits:

It was observed that at 0.2mg/mL, rot pathogens were inhibited thus: *Aspergillus spp* (21.00mm), *Cladosporium spp* (19.00mm) and *Fusarium solani* (13.00mm), *Rhizopus stolonifer* (17.00mm), *Mucor spp* (17.00mm), *Alternaria sp* (15.00mm) (Table 2).

Table 2: Antifungal activity of clove (*Syzygium aromaticum*) methanol extract on fungal pathogens of green pepper.

| Isolates | Diameter of zones of inhibition (mm) | | | | |
|----------------------------|--------------------------------------|--------------------|--------------------|--------------------|-------------------|
| | 0.2 | 0.4 | 0.6 | 0.8 | 1.0 |
| | Concentrations (mg/mL) | | | | |
| <i>Aspergillus spp</i> | 21.00 ^a | 18.00 ^a | 14.00 ^a | 10.00 ^a | 8.00 ^a |
| <i>Cladosporium spp</i> | 19.00 ^b | 15.00 ^b | 12.00 ^c | 9.00 ^b | 7.00 ^b |
| <i>Fusarium solani</i> | 13.00 ^c | 10.00 ^d | 8.00 ^e | 0.00 ^d | 0.00 ^d |
| <i>Rhizopus stolonifer</i> | 17.00 ^c | 15.00 ^b | 13.00 ^b | 10.00 ^a | 8.00 ^a |
| <i>Mucor spp</i> | 17.00 ^c | 15.00 ^b | 13.00 ^b | 10.00 ^a | 8.00 ^a |
| <i>Alternaria spp</i> | 15.00 ^d | 13.00 ^c | 10.00 ^d | 8.00 ^c | 6.00 ^c |

Values are mean \pm standard error of the mean for bio-assay conducted in triplicate. Means followed by the same letter(s) are not significantly different (multivariate analysis, Fisher's protected LSD at $p \leq 0.05$).

Key: Control Nil.

The antifungal effects of methanolic extract of ginger (*Zingiber officinale*).

It was observed that at 1.0mg/mL, rot pathogens were inhibited thus: *Cladosporium spp* (15.00mm), *Rhizopus stolonifer* (17.00mm), *Fusarium solani* (13.00mm), *Aspergillus spp* (18.00mm), *Cladosporium spp* (15.00mm) (Table 3).

Table 3. Antifungal activity of ginger (*Zingiber officinale*) methanol extract against fungi pathogen of green pepper.

| Isolates | Diameter of zones of inhibition (mm) | | | | |
|----------------------------|--------------------------------------|--------------------|--------------------|--------------------|-------------------|
| | 1.0 | 0.8 | 0.6 | 0.4 | 0.2 |
| | Concentrations (mg/mL) | | | | |
| <i>Aspergillus spp</i> | 18.00 ^b | 0.00 ^d | 0.00 ^d | 0.00 ^d | 0.00 ^c |
| <i>Cladosporium spp</i> | 15.00 ^d | 12.00 ^b | 10.00 ^b | 8.00 ^b | 7.00 ^b |
| <i>Fusarium solani</i> | 13.00 ^f | 10.00 ^c | 8.00 ^c | 7.00 ^c | 0.00 ^c |
| <i>Rhizopus stolonifer</i> | 17.00 ^c | 15.00 ^a | 13.00 ^a | 10.00 ^a | 8.00 ^a |
| <i>Mucor spp</i> | 14.00 ^c | 12.00 ^b | 10.00 ^b | 8.00 ^b | 0.00 ^c |
| <i>Aiternaria spp</i> | 20.00 ^a | 10.00 ^c | 10.00 ^b | 8.00 ^b | 8.00 ^a |

Values are mean \pm standard error of the mean for bio assay conducted in triplicate. Means followed by the same letter(s) are not significantly different (multivariate analysis, Fisher's protected LSD at $p \leq 0.05$).

Key: Control Nil.

The result of garlic (*Allium sativum*) methanolic extract against fungal isolates of green pepper fruits at 0.1mg/mL.

It shows that *Aspergillus spp* (12.00mm), *Cladosporium spp* (24.00mm), *Fusarium solani* (25.70mm) *Rhizopus stolonifer* (20.00mm), *Mucor spp* (22.00mm), *Alternaria spp* (19.00mm), *Fusarium solani* (25.70mm) and *Cladosporium spp* (24.00mm) (Table 4).

Table 4: Antifungal activity of garlic methanol extract on fungi pathogen of green pepper.

| Isolates | Diameter of zones of inhibition (mm) | | | | |
|----------------------------|--------------------------------------|--------|--------|-------|-------|
| | 1.0 | 0.8 | 0.6 | 0.4 | 0.2 |
| | Concentrations (mg/mL) | | | | |
| <i>Aspergillus spp</i> | 20.00d | 10.00f | 9.00d | 0.00b | 0.00a |
| <i>Cladosporium spp</i> | 24.00b | 12.70b | 9.50c | 0.00b | 0.00a |
| <i>Fusarium solani</i> | 25.70a | 15.20a | 8.50e | 8.00a | 0.00a |
| <i>Rhizopus stolonifer</i> | 20.00d | 12.00d | 8.00f | 0.00b | 0.00a |
| <i>Mucor spp</i> | 22.00c | 11.90e | 10.80a | 0.00b | 0.00a |
| <i>Alternaria spp</i> | 19.00e | 13.10d | 10.30b | 8.00a | 6.00a |

Values are mean \pm standard error of the mean for bio-assay conducted in triplicate. Means followed by the same letter(s) are not significantly different (multivariate analysis, Fisher's protected LSD at $p \leq 0.05$). Key: Control Nil.

The antifungal effect of methanolic extract of various spices against fungi of green pepper.

Antifungal effect of garlic was most effective against *Cladosporium sp* (24), *Fusarium sp* (25), *Rhizopus sp* (20), *Mucor* (22) while clove was most active against *Aspergillus sp* (21mm) (Table 5).

Table 5. Antifungal activity of methanol extract of different spices against isolated fungi.

| Isolates | Fluconazole(control) mm | Clove (100 μ l)mm | Ginger (100 μ l)mm | Garlic (100 μ l) mm |
|----------------------------|-------------------------|-----------------------|------------------------|-------------------------|
| <i>Aspergillus spp</i> | 22 | 21 | 18 | 20 |
| <i>Cladosporium spp</i> | 18 | 19 | 15 | 24 |
| <i>Fusarium solani</i> | 19 | 13 | 13 | 25 |
| <i>Rhizopus stolonifer</i> | 22 | 17 | 17 | 20 |
| <i>Mucor spp</i> | 28 | 17 | 14 | 22 |
| <i>Alternaria spp</i> | 29 | 15 | 20 | 19 |

Proximate composition of infected green pepper fruits with various fungi

Moisture content of *Cladosporium spp* (19.68) infected green pepper was highest, total ash was highest in *Mucor sp* infected green pepper (18.87), crude fibre was highest in green pepper infected with

Alternaria spp (23.98), crude fat was highest in green pepper infected with *Mucor sp* and *Fusarium solani* (9.97), crude protein was highest in green pepper infected with *Cladosporium spp* (15.96), carbohydrate was highest in green pepper infected with *Aspergillus spp* (32.85) (Table 6).

Table 6. Proximate composition of infected green pepper fruits with various fungi.

| Parameters (%) | Sample codes | | | | | | |
|------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| | Control | A | B | C | D | E | F |
| Moisture content | 18.41 ^c | 18.59 ^d | 19.68 ^c | 19.24 ^c | 19.08 ^c | 19.26 ^c | 19.20 ^c |
| Total ash | 18.41 ^c | 18.61 ^c | 18.81 ^d | 18.85 ^d | 18.77 ^d | 18.87 ^d | 18.71 ^d |
| Crude fiber | 22.90 ^b | 23.31 ^b | 23.22 ^b | 23.86 ^b | 23.52 ^b | 23.21 ^b | 23.98 ^b |
| Crude fat | 9.25 ^c | 9.35 ^f | 9.92 ^f | 9.97 ^f | 9.91 ^f | 9.97 ^f | 9.95 ^f |
| Crude protein | 15.89 ^d | 15.63 ^e | 15.96 ^e | 15.16 ^e | 15.67 ^e | 15.68 ^e | 15.01 ^e |
| Carbohydrate | 33.75 ^a | 32.85 ^a | 33.57 ^a | 33.12 ^a | 33.22 ^a | 32.78 ^a | 33.57 ^a |

Values are mean \pm standard error of the mean for bio-assay conducted in triplicate. Means followed by the same letter are not significantly different (multivariate analysis, Fisher's protected LSD at $p \leq 0.05$). Harmonic mean sample size 6.00.

Key:

A = *Aspergillus spp*, B = *Cladosporium spp*, C = *Fusarium solani*, D = *Rhizopus stolonifera*, E = *Mucor spp*, F = *Alternaria spp*.

DISCUSSION

The findings of this study showed that *Aspergillus spp* with highest percentage occurrence of 36%, *Alternaria spp*. (28%), *Cladosporium spp*, (16%), *Fusarium solani* (8%), *Mucor spp*(8%) and *Rhizopus stolonifer* (4%) were post harvest rot pathogens of pepper fruits sold in major market in Ado Ekiti metropolis, Nigeria. A number of these microbes were isolated from deteriorated mango and oranges by Ijato, *et al* (2021). Similarly, Ugwu *et al.* (2014) isolated *Candida tropicalis*, *Penicillium notatum*, *Aspergillus niger*, *Fusarium oxysporum*, *Absidia corymbifera* and *Rhizopus stolonifer* from post-harvest pepper fruits, this was similar to the result obtained in their study. Also, the isolated fungi in this study agreed with fungal species associated with post-harvest rot of common fruits pepper in Sokoto metropolis, Nigeria as reported by Salau, (2012). Occurrence of *Aspergillus niger*, *Fusarium* species and *Mucor* species as identified in this study agrees with the work of Mensah and Owusu (2011) that isolated *Aspergillus niger*, *Fusarium* species and *Mucor* species from *Capsicum annum*, *Abelmoscus esculentus*, and *Lycopersicon esculentum* in Accra metropolis.

However, *Aspergillus niger* was found to be the most abundant fungus of pepper fruit in this study, this is similar to the report of Mensah and Owusu, (2011) that listed *Aspergillus niger* as one of the most common fungal species found on fruits in Accra metropolis. It also agrees with Chiejina (2008) who indicated that *Aspergillus* was isolated from 79.5% of the samples. Several studies have also reported that *Aspergillus spp*. are associated with spoilage of mango, tomatoes, pepper, apricot, orange, lemon, peach, apple, kiwi,

mango and banana (Rashad *et al.*, 2011, Ijato *et al.*, 2022 a & b). Onuorah and Orji (2015) reported that *Aspergillus* had the highest decay diameter among other fungi associated with pepper spoilage. The fungal isolates from this study have been reported to form mycotoxins (Onuorah and Orji, 2015). Generally, fungi that cause spoilage are considered toxigenic or pathogenic (Al-Hindi *et al.*, 2011). Also, some of the fungi isolated in this study have been reported to produce secondary metabolites in plants tissues which are harmful to humans and animals (Baiyewu *et al.*, 2007).

In this study, garlic was most effective compared to other spices used. Garlic has highest inhibitory effect on *Cladosporium spp* (24 mm), *Fusarium solani* (25 mm), *Rhizopus stolonifer* (20 mm) and *Mucor spp* (22 mm). several *in vitro* studies have shown the efficacy of certain medicinal herbs or plant extracts against *Colletotrichum spp* (Saravanakumar *et al.*, 2011; Ajith *et al.*, 2012). Also, Alves *et al.* (2015) reported the efficacy of 1% aqueous or 20% ethanol plant extracts to control bell pepper (*C. annum*) anthracnose caused by *C. acutatum* after the fruits were treated with plant extracts before inoculation with *C. acutatum* conidial suspension of *C. acutatum* (2×10^5) conidia/ml. Garlic is more effective in this study as compared to commercial antibiotics. Ijato, 2021 a, b, c & d reported the used of plant extracts as an alternative remedy for the control of various plant pathogens ranging from fungi to bacteria. Garlic is not only effective against gram-positive and gram-negative bacteria but also possess antiviral and antifungal activities (Benavides *et al.*, 2007). These studies elaborate the effective antimicrobial quality of different spices especially garlic. Therefore, garlic can be used as an effective anti fungal agent against many fungal diseases and also used as natural food preservative.

CONFLICT OF INTEREST STATEMENT

There is no conflict of interest in this research as all the authors agreed on the procedures and results presented in this research.

REFERENCES

- [1]. Ademoh, O. F., Afolabi, A. M., Orisasona B. A. and Olowolaju E. D. (2017). Isolation and Identification of Rot Fungi on Post-Harvest of Pepper (*Capsicum annum* L.) Fruits. *AASCIT Journal of Biology*. 3(5): 24-29.
- [2]. Al-Hindi, R. R., Al-Najada, A. R. and Mohamed, S. A. (2011) Isolation and identification of some fruit spoilage fungi: Screening of plant cell wall degrading enzymes. *African Journal of Microbiology Research*. 5(4): 443-448.
- [3]. Alves K. F., Laranjeira D. and Câmara, M. P. S (2015). Efficacy of plant extracts for anthracnose control in bell pepper fruits under controlled conditions. *Horticultura Brasileira* 33(3):332-338
- [4]. AOAC (2000). Official methods of analysis, 17th Edition. Association of Official Analytical Chemists, Arlington, Virginia.
- [5]. AOAC (2009). Official methods of analysis. 21st Edition. Association of Official Analytical Chemists, Arlington, Virginia.
- [6]. Baiyewu, R.A., Amusa, N. A., Ayoola, O.A. and Babalola, O.O. (2007). Survey of the post-harvest diseases and aflatoxin contamination of marketed pawpaw fruit (*Carica papaya* L) in South-Western Nigeria. *African Journal of Agricultural Research*. 2(4): 178-181
- [7]. Balogun, O.S., Odeyemi, G.A. and Fawole, O.B. (2005) Evaluation of the pathogenic effect of some fungal isolates on fruits and seedlings of pepper (*Capsicum* species). *Journal of Agricultural Research and Development* 4 (2), 159 – 169.
- [8]. Benavides, G. A., Squadrito, G.L, Mills R.W., Patel H.D. and Isbell T.S (2007). Hydrogen sulfide mediates the vasoactivity of garlic. *Proc Natl Acad Sci*. 104(46): 17977-17982.
- [9]. Chandrashekhara, C., Kumar R., Bhatt, J.C. and Chandrashekhara, K.N (2012). Suppressive soils in plant disease management. In: Singh, V.K., Singh Y, Singh A (eds) Eco-friendly innovative approaches in plant disease management. International Book Distributors, India, pp 241–256.
- [10]. Cheesbrough, M. (2000) District Laboratory Practice Manual in Tropical Countries Part 2. Cambridge University Press, Cambridge, 178-179.
- [11]. Chiejina, N. V. (2008). Mycoflora of some salad vegetables. *Biological Research*. 6(2): 392-39.
- [12]. Fagbohun, E. D. and Bamikole, A. M. (2019). Antifungal Effects of Methanolic Extract of Stem Bark of *Bridelia ferruginea* Benth. Leaves of *Aloe vera* L. and Stem Bark of *Alstonia boonei* De Wild. *Microbiology Research Journal International*. 27(2): 1-11
- [13]. Fawole, M.O. and Oso, B.A. (1995) Laboratory manual of microbiology. Spectrum Books Limited Ibadan, Owerri, 71-81.
- [14]. Heikki H., Melita Z., Gun-Britt H., Ingeborg Menzler-Hokkanen, W Buchs, Z Klukowski, Luik A., Nilsson C., Ulber, B. and Williams, I.H. (2006). Effectiveness of entomopathogenic nematodes in the control of oilseed rape pests. International symposium on integrated pest management in oilseed rape, 3rd-5th April 2006, Paulinerkirche, Göttingen
- [15]. Heiser, C. B., Mason, J. R. and Pickersgill, B. (2010). Names for the cultivated *Capsicum* Species (Solanaceae) Taxon, 18 (3): 277-283.
- [16]. Ijato J. Y, Akinjogunla O. Joseph, Divine-Anthony Ofon-mbuk(2021). *Aspergillus flavus*, *Rhizopus stolonifer* and *Mucor* spp. Associated with Deteriorated Mango and Orange Fruits: Occurrence and In vitro Susceptibility to Extracts of *Aspilia*. *Tropical Journal of Natural Product Research* 5 (9): 1650-1655
- [17]. Ijato J. Y(2021a). Evaluation of bio-protective capacity of some botanicals against post harvest fruit rot microbes of pepper (*Capsicum annum* L). *Researcher* 13 (3):14-20
- [18]. Ijato J. Y(2021b). Bioassay of some plant oil against bacterial rot pathogens. *Am Sci j*. 17 (5), 30-36
- [19]. Ijato J. Y(2021c). Antibacterial Activity of *Vernonia Amygdalina* on Post Harvest Organisms Associated with Cocoyam (*Colocasia Esculentus* L) Corms Rot. *J. Pharm Microbiol* 7(2):1-4
- [20]. Ijato J. Y(2021d). Assessment of antibacterial capacity of *Vernonia amygdalina* against post harvest fruit rot organisms of okra (*Abelmoshus esculentus* L moench). *Life Science Journal* 18 (5): 50-53
- [21]. Ijato J. Y, Olajide, O. O. and Ojo (2022a). Studies On Bioactivities Of Various Parts Of *Murraya Koenigii* (L) Spreng (curry Tree) On Fungal Isolates From Tomatoes. *advanced Journal of Plant Biology*. 3 (1):1-6
- [22]. Ijato J. Y, Olajide, O. O. and Ojo (2022b). *Rhizoctonia solani*, *Aspergillus niger*, *Streptococcus pyrogenes*, *Alcaligenes faecalis* and *Proteus vulgaris* selectively associated with two varieties of banana and effects of storage conditions on nutritional composition

- of banana. *International Journal of Frontline Research and Reviews* 1 (1): 007–013
- [23]. Mensah, K. and Owusu, E. (2011). Fruit bornemicroflora of *Capsicum annum* L. (pepper) *Abelmoschus esculentus* L. Moench (Okra), and *Lycopersicon esculentum* Mill. (tomato) from Accra metropolis. *African Journal of food Sciences* (1): 1-7.
- [24]. Naresh Magan and David Aldred (2007). Post-harvest control strategies: minimizing mycotoxins in the food chain. *Int J Food Microbiol.* 119(1-2):131-9.
- [25]. Nene, Z.H, Thapilyal (2002). Management of mushroom pathogens through botanicals. *Ind. Phytopathol.*, 58:189-193.
- [26]. Onuorah, S. and Orji, M.U. (2015). Fungi associated with the spoilage of post-harvest tomato fruits sold in major markets in Awka, Nigeria. *Universal Journal of Microbiology Research.* 3(2):11-16.
- [27]. Parveen, S., Gupta, D.B, Dass S., Kumar, A., Pandey, A., Chakraborty, S. and Chakraborty, N. (2016). Chickpea Ferritin CaFer1 Participates in Oxidative Stress Response, and Promotes Growth and Development. *Sci Rep.*, 6:31218
- [28]. Rashad al-hindi, A., Rashed A., and Saleh, A. H. M. (2011). Isolation and identification of some fruit spoilage fungi: Screening of plant cell wall degrading enzymes. *African Journal of Microbiology Research* 5(4):443-448
- [29]. Salau, I. A. (2012) Studies of fungi associated with human skin and vegetable disease in fadamaland of Sokoto metropolis, Sokoto state, Nigeria. M.Sc Thesis, Usman Dan Fodio University Sokoto, Nigeria.
- [30]. Saravanakumar, M., Kavino, T., Raguchander, P., Subbian R., Samiyappan (2011). Plant growth promoting bacteria enhance water stress resistance in green gram plants. *Acta Physiol Plant* , 33:203–209
- [31]. Sasvari, J. (2005). Paprika: A Spicy Memoir from Hungary. Toronto, ON: Can West Books. p202.
- [32]. Snowdon, A. (1990). A Colour Atlas of Post-Harvest Disorders and Diseases of Fruits and Vegetables, Vol. 1 Pome Fruits 302 pp
- [33]. Sohail, A. Alsohaili and Bayan M. Bani-Hasan (2018). Morphological and Molecular Identification of Fungi Isolated from Different Environmental Sources in the Northern Eastern Desert of Jordan. *Jordan Journal of Biological Sciences.* 11 (3): 329 - 337
- [34]. Ugwu, C.O., Chukwuezi, F.O, Ozougwu, V.C.O. (2014) Microbial agents of tomato spoilage in Onitsha metropolis. *Advances in Biological Research.* 8(2):87-93.
- [35]. Zorya, S., Morgan, N., Diaz Rios, L., Hodges, R., Bennett, B., Stathers, T. and Lamb, J. (2011). Missing Food, the Case of Post-harvest Grain Losses in Sub-Saharan Africa. Report No. 60371. The World Bank, Washington DC. 10pp.

8/24/2022